

On Weakly Coherent Rings

Chahrazade Bakkari and Najib Mahdou

Department of Mathematics, Faculty of Science and Technology of Fez,
Box 2202, University S. M. Ben Abdellah Fez, Morocco
cbakkari@hotmail.com
mahdou@hotmail.com

Abstract. In this paper, we define weakly coherent rings, and examine the transfer of these rings to homomorphic image, trivial ring extension, localization, and direct product. These results provide examples of weakly coherent rings that are not coherent rings. We show that the class of weakly coherent rings is not stable by localization. Also, we show that the class of weakly coherent rings and the class of strongly 2-coherent rings are not comparable.

Key Words. Weakly coherent ring, coherent ring, homomorphic image, trivial ring extension, localization, direct product.

2000 Mathematics Subject Classification: 13F05, 13B05, 13A15, 13D05, 13B25.

1 Introduction

Throughout this paper all rings are assumed to be commutative with identity elements and all modules are unital.

Let R be a commutative ring. We say that an ideal is regular if it contains a regular element, i.e; a non-zerodivisor element.

For a nonnegative integer n , an R -module E is n -presented if there is an exact sequence of R -modules:

$$F_n \rightarrow F_{n-1} \rightarrow \dots F_1 \rightarrow F_0 \rightarrow E \rightarrow 0$$

where each F_i is a finitely generated free R -module. In particular, 0-presented and 1-presented R -modules are, respectively, finitely generated and finitely presented R -modules.

A ring R is coherent if every finitely generated ideal of R is finitely presented; equivalently, if $(0 : a)$ and $I \cap J$ are finitely generated for every $a \in R$ and any two finitely generated ideals I and J of R [5]. Examples of coherent rings are Noetherian rings, Boolean algebras, von Neumann regular rings, valuation rings, and Prüfer/semihereditary rings. See for instance [5].

In this paper, we investigate a particular class of coherent rings that we call weakly coherent rings. A ring R is called a weakly coherent ring if for each proper ideals $I \subseteq J$ such that I is finitely generated and J is finitely presented, then I is finitely presented. If R is coherent, then R is naturally weak coherent. Our aim in this paper is to prove that the converse is false in general.

We say that R is strong n -coherent if each n -presented R -module is $(n + 1)$ -presented. In particular, any coherent ring (i.e., 1-coherent ring) is a strong 2-coherent ring. This led us to consider the relation between the class of weakly coherent rings and the class of strong 2-coherent rings.

Let A be a ring, E be an A -module and $R := A \ltimes E$ be the set of pairs (a, e) with pairwise addition and multiplication given by $(a, e)(b, f) = (ab, af + be)$. R is called the trivial ring extension of A by E . Considerable work has been concerned with trivial ring extensions. Part of it has been summarized in Glaz's book [5]) and Huckaba's book (where R is called the idealization of E by A) [7].

In the context of non-total ring of quotients (i.e., ring containing a regular element), we show that the notion of weak coherent coincide with the definition of coherent ring. The goal of this work is to exhibit a class of non-coherent weakly coherent rings. We show that the class of weakly coherent rings is not stable by localization. Also, we show that the class of weakly coherent rings and the class of strong 2-coherent rings are not comparable. For this purpose, we study the transfer of this property to homomorphic image, trivial ring extension, and direct product.

2 Main Results

Recall that for nonnegative integers n and d , we say that a ring R is an (n, d) -ring if $pd_R(E) \leq d$ for each n -presented R -module E (as usual, $pd_R E$ denotes the projective dimension of E as an R -module). See for instance [2, 8, 9, 10, 11].

We begin this section by giving an example of non-coherent weakly coherent ring.

Example 2.1 *Let (A, M) be a local ring with maximal ideal M , E be an A/M -vector space with infinite rank, and let $R := A \ltimes E$ be the trivial ring extension of A by E . Then:*

- 1) *R is a weak coherent ring.*
- 2) *R is not a coherent ring. $\square$*

Proof. 1) It suffices to show that there is no finitely presented proper ideal J of R . Deny. Let J be a finitely presented proper ideal of R . Then J is free since R is a local $(2, 0)$ -ring by [11, Theorem 2.1(1)], that is $J = Ra$ for some regular element a of R ; a contradiction since $J \subseteq M \ltimes E$ and $(M \ltimes E)(0, e) = (0, 0)$ for each $e \in E - \{0\}$. Hence R is a weak coherent ring.

2) We claim that R is not coherent. Deny. Assume that R is coherent. But, R is a $(2, 0)$ -ring by [11, Theorem 2.1(1)]. Hence, R is Von Neumann regular ring since it is coherent, a contradiction by [11, Theorem 2.1(2)]. Hence, R is not coherent, as desired. $\square$

Now, we give a sufficient condition to have equivalence between a coherence and weakly coherence properties.

Proposition 2.2 *Let R be a commutative ring, then:*

- 1) *If R is a coherent ring, then R is a weak coherent ring.*
- 2) *Assume that R contains a regular element (that is R is not a total ring of quotients). Then R is a coherent ring if and only if R is a weak coherent ring. $\square$*

Proof. 1) Clear.

2) It remains to show that if R is a weak coherent ring and contains a regular element a , then R is a coherent ring. Let I be a finitely generated proper ideal of R . Hence, $aI \subseteq aR$, aI is a finitely generated proper ideal of R , and $aR(\cong R)$ is a finitely presented proper ideal of R . Therefore, aI is a finitely presented ideal of R and so $I(\cong aI)$ (since a is regular) is finitely presented, as desired. $\square$

Remark 2.3 *By the above result, a non-coherent weak coherent ring is necessary a total ring of quotients. $\square$*

Now, we investigate the homomorphic image of weak coherent rings.

Theorem 2.4 *Let R be a weak coherent ring and I be a finitely generated ideal of R . Then R/I is a weak coherent ring.*

Proof. Let $L/I \subseteq J/I$ be two finitely generated proper ideals of R/I such that J/I is finitely presented. Our aim is to show that L/I is finitely presented.

Remark that $L \subseteq J$ are two finitely generated proper ideals of R . We claim that J is finitely presented.

Indeed, there exists an exacte sequence of (R/I) -modules:

$$0 \rightarrow T \rightarrow (R/I)^n \rightarrow J/I \rightarrow 0(*)$$

where n is a positive integer and T is a finitely generated (R/I) -module (since J/I is a finitely presented ideal of R/I). Hence, T is a finitely generated R -module. On the other hand, R/I is a finitely presented R -module (since I is a finitely generated ideal of R and by the exact sequence of R -modules $0 \rightarrow I \rightarrow R \rightarrow R/I \rightarrow 0$). Therefore, J/I is a finitely presented R -module by exact sequence $(*)$ considered as an exact sequence of R -modules. Consequently, J is a finitely presented ideal of R by the exact sequence of R -modules $0 \rightarrow I \rightarrow J \rightarrow J/I \rightarrow 0$, as desired.

Now, we have $L \subseteq J$, L is a finitely generated ideal, and J is a finitely presented ideal; so L is a finitely presented ideal of R since R is a weakly coherent ring. Hence, the exact sequence of R -modules $0 \rightarrow I \rightarrow L \rightarrow L/I \rightarrow 0$ shows that L/I is a finitely presented R -module. We claim that L/I is a finitely presented ideal of R/I and this completes the proof of Theorem 2.4.

Indeed, since L/I is a finitely generated ideal of R/I , consider the exact sequence of (R/I) -modules:

$$0 \rightarrow S \rightarrow (R/I)^m \rightarrow L/I \rightarrow 0(**)$$

where m is a positive integer and S is an (R/I) -module. The exact sequence $(**)$ is also an exact sequence of R -modules; hence S is a finitely generated R -module since L/I and R/I are finitely presented R -modules. Therefore, S is also a finitely

generated (R/I) -module and the exact sequence of (R/I) -modules $(**)$ shows that L/I is a finitely presented ideal of R/I , as desired. $\square$

The condition " I is a finitely generated ideal of R " is necessary in Theorem 2.4 as the following example shows:

Example 2.5 *Let (A, M) be a non-coherent local domain with maximal ideal M , E be an A/M -vector space with infinite rank, let $R := A \ltimes E$ be the trivial ring extension of A by E , and set $I := 0 \ltimes E$. Then:*

- 1) *R is a weak coherent ring by Example 2.1.*
- 2) *$R/I(\cong A)$ is not a weak coherent ring (by Proposition 2.2 since A is not a coherent domain). $\square$*

Now, we investigate a weak coherent property in a particular class of total rings of quotients; namely, those arising as trivial ring extensions of local rings by vector spaces over the residue fields. The following main result enriches the literature with original examples of non-coherent weak coherent rings.

Theorem 2.6 *Let (A, M) be a local ring with maximal ideal M , E be an A -module such that $ME = 0$, and let $R := A \ltimes E$ be the trivial ring extension of A by E . Then R is a weak coherent ring if and only if one of the following two properties holds:*

- 1) *E is an A/M -vector space with infinite rank.*
- 2) *E is an A/M -vector space with finite rank and A is weak coherent. $\square$*

Proof. Assume that R is a weak coherent ring and E is an A/M -vector space with finite rank. Our aim is to show that A is weak coherent.

Let $I \subseteq J$ be two proper ideals of A such that I is finitely generated and J is finitely presented. Then $(I \ltimes 0) \subseteq (J \ltimes 0)$ are two finitely generated proper ideals of R . We claim that $J \ltimes 0$ is a finitely presented ideal of R .

Indeed, let $J := \sum_{i=1}^n Aa_i$ for some positive integer n and some $a_i \in J$, and consider the exact sequence of A -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow A^n \xrightarrow{u} J \rightarrow 0$$

where u is defined by $u((b_i)_{1 \leq i \leq n}) = \sum_{i=1}^n b_i a_i$. Hence, $\text{Ker}(u)(: \{(b_i)_{1 \leq i \leq n} \in A^n / \sum_{i=1}^n a_i b_i = 0\})$ is a finitely generated A -module (since J is a finitely presented

ideal of A). Now, consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(v) \rightarrow R^n \xrightarrow{v} J \rtimes 0 \rightarrow 0$$

where $v((b_i, e_i)_{1 \leq i \leq n}) = \sum_{i=1}^n (b_i, e_i)(a_i, 0)$. But,

$$\begin{aligned} \text{Ker}(v) &= \{(b_i, e_i)_{1 \leq i \leq n} \in R^n / \sum_{i=1}^n (a_i, 0)(b_i, e_i) = 0\} \\ &= \{(b_i, e_i)_{1 \leq i \leq n} \in R^n / \sum_{i=1}^n a_i b_i = 0\} \\ &= \text{Ker}(u) \rtimes E^n \end{aligned}$$

(since $a_i \in J \subseteq M$) which is finitely generated R -module (since $\text{Ker}(u)$ and E are finitely generated A -modules). Hence, $J \rtimes 0$ is a finitely presented (proper) ideal of R and so $(I \rtimes 0) (\subseteq J \rtimes 0)$ is a finitely presented ideal of R since R is a weak coherent ring. Therefore, by the same reasoning as for J above, we can show that I is finitely presented and this shows that A is a weak coherent ring.

Conversely, if $\dim_{A/M} E = \infty$, then R is a weak coherent ring by Example 2.1. Now, assume that $\dim_{A/M} E < \infty$ and A is a weak coherent ring and our aim is to show that R is a weak coherent ring.

Let $I(= \sum_{i=1}^n R(a_i, e_i)) \subseteq J(= \sum_{i=1}^m R(b_i, f_i))$ be two proper ideals of R such that n, m are positive integers, $a_i, b_j \in A$ and $e_i, f_j \in E$ for each i, j , and J is finitely presented. We wish to show that I is finitely presented. Two cases are then possible:

Case 1. $b_i = 0$ for all $i = 1, \dots, m$.

In this case, $a_i = 0$ for all $i = 1, \dots, n$ and $I := 0 \rtimes E_1$ and $J := 0 \rtimes E_2$ for some (A/M) -vector subspace E_1 and E_2 of E . Assume that $(e_i)_{i=1, \dots, n}$ and $(f_i)_{i=1, \dots, m}$ are respectively basis of the (A/M) -vector space E_1 and E_2 . Consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow R^m \xrightarrow{u} J(= 0 \rtimes E_2) \rightarrow 0$$

where $u((c_i, g_i)_{1 \leq i \leq m}) = \sum_{i=1}^m (c_i, g_i)(0, f_i) = (0, \sum_{i=1}^m c_i f_i)$. Hence, $\text{Ker}(u) = M^m \rtimes E^m (= (M \rtimes E)^m)$ since $(f_i)_{i=1, \dots, m}$ is a basis of the (A/M) -vector space E_2 and so M is a finitely generated ideal of A (since J is a finitely presented ideal of R). Therefore, the exact sequence of R -modules:

$$0 \rightarrow M^n \rtimes E^n \rightarrow R^n \xrightarrow{v} I(= 0 \rtimes E_1) \rightarrow 0$$

where $v((c_i, g_i)_{1 \leq i \leq n}) = \sum_{i=1}^n (c_i, g_i)(0, e_i)$ shows that I is a finitely presented ideal of R (since $M^n \rtimes E^n$ is a finitely generated R -module), as desired.

Case 2. $b_i \neq 0$ for some $i = 1, \dots, m$.

We may assume that $((a_i, e_i)_{i=1, \dots, n})$ and $((b_i, f_i)_{i=1, \dots, m})$ are minimal generating sets respectively of I and J . Set $I_0 := \sum_{i=1}^n Aa_i$ and $J_0 := \sum_{i=1}^m Ab_i$. Consider the exact sequence of R -modules:

$$0 \rightarrow \text{Ker}(u) \rightarrow R^m \xrightarrow{u} J \rightarrow 0$$

where $u((c_i, g_i)_{1 \leq i \leq m}) = \sum_{i=1}^m (c_i, g_i)(b_i, f_i)$. But $\text{Ker}(u) \subseteq (M \rtimes E)^m$ by [12, Lemma 4.43, p.134]. Hence,

$$\begin{aligned} \text{Ker}(u) &= \{(c_i, g_i)_{1 \leq i \leq m} \in R^m / \sum_{i=1}^m (c_i, g_i)(b_i, f_i) = 0\} \\ &= \{(c_i, g_i)_{1 \leq i \leq m} \in R^m / \sum_{i=1}^m c_i b_i = 0\} \\ &= V \rtimes E^m \end{aligned}$$

where $V := \{(c_i)_{1 \leq i \leq m} \in A^m / \sum_{i=1}^m c_i b_i = 0\}$. Also, we have the exact sequence of R -modules:

$$0 \rightarrow V \rtimes E^m \rightarrow R^m \xrightarrow{w} J_0 \rtimes 0 \rightarrow 0$$

where $w((c_i, g_i)_{1 \leq i \leq m}) = \sum_{i=1}^m (c_i, g_i)(b_i, 0)$. Therefore, we may assume that $J = J_0 \rtimes 0$ and so $I = I_0 \rtimes 0$ (since $I \subseteq J$). On the other hand, V is a finitely generated A -module since $V \rtimes E^m (= \text{Ker}(u))$ is a finitely generated R -module (by the above exact sequence since J is finitely presented) and so J_0 is a finitely presented ideal of A by the exact sequence of A -modules:

$$0 \rightarrow V \rightarrow A^m \xrightarrow{v} J_0 \rightarrow 0$$

where $v((c_i)_{1 \leq i \leq m}) = \sum_{i=1}^m c_i b_i$. Hence, I_0 is a finitely presented ideal of A since A is weak coherent and $I_0 \subseteq J_0$. Therefore, by the same reasoning as for J , we may show that I is a finitely presented ideal of R and this completes the proof of Theorem 2.6. $\square$

Now, we are able to give a new class of a non-coherent weak coherent rings.

Example 2.7 Let (A, M) be a local coherent ring with non-finitely generated maximal ideal M , E an A -module such that $ME = 0$, let $R := A \ltimes E$ be the trivial ring extension of A by E , and set $I := 0 \ltimes E$. Then:

- 1) R is a weak coherent ring by Theorem 2.6.
 - 2) R is not a coherent ring by [9, Theorem 2.6(2)] since M is not finitely generated.
-

The localization of a weak coherent ring is not always a weak coherent ring as the following Example shows:

Example 2.8 Let $A := \mathbb{Z}_2 + X\mathbb{R}[[X]]$ a local ring with maximal ideal $M = 2\mathbb{Z}_2 + X\mathbb{R}[[X]]$, E be an A/M -vector space with infinite rank and let $R := A \ltimes E$ be the trivial ring extension of A by E . Let S be the multiplicative subset of R given by $S := \{(2, 0)^n \mid n \in \mathbb{N}\}$ and S_0 the multiplicative subset of A given by $S_0 := \{2^n \mid n \in \mathbb{N}\}$. Then:

- 1) R is a weak coherent ring.
- 2) $S^{-1}R$ is not a weak coherent ring. □

Proof. 1) Clear by Example 2.1.

2) Since $2E \subseteq ME = 0$ and $2 \in S_0$, then $S_0^{-1}E = 0$. Thus, $S^{-1}(0 \ltimes E) = 0$ and so $S^{-1}R = \{ \frac{(a, 0)}{(s, 0)} \mid a \in A \text{ and } s \in S_0 \}$ which is clearly isomorphic to a ring $S_0^{-1}A$. But $S_0^{-1}A = S_0^{-1}\mathbb{Z}_2 + X\mathbb{R}[[X]] = \mathbb{Q} + X\mathbb{R}[[X]]$ which is not a coherent domain by [5, Theorem 5.2.3]. Therefore, $S^{-1}R$ is not a weak coherent ring by Proposition 2.2, as desired. □

We know that a coherent ring is weak coherent and strong 2-coherent. The following two examples show that the class of weakly finite conductor rings and the class of 2-coherent rings are not comparable.

Example 2.9 Let R be a non-coherent strong 2-coherent domain (see for example [10, Theorem 3.1]). Then, R is not a weak coherent domain (by Proposition 2.2 since R is not a coherent domain). □

Example 2.10 Let (A, M) be a local coherent domain with non-finitely generated maximal ideal M , and let $R := A \ltimes (A/M)$ be the trivial ring extension of A by A/M . Then:

- 1) R is a weak coherent ring by Theorem 2.6.
- 2) R is not a strong 2-coherent ring by [8, Theorem 3.1]. $\square$

Next, we study the transfer of weak coherent property to direct products.

Proposition 2.11 Let $(R_i)_{i=1,\dots,n}$ be a family of rings. Then, $\prod_{i=1}^n R_i$ is a weak coherent ring if and only if R_i is a weak coherent ring for each $i = 1, \dots, n$. $\square$

We need the following Lemma before proving Proposition 2.11.

Lemma 2.12 ([10, Lemma 2.5(1)]) Let $(R_i)_{i=1,2}$ be a family of rings and E_i an R_i -module for $i = 1, 2$. Then $E_1 \amalg E_2$ is a finitely generated (resp., finitely presented) $R_1 \amalg R_2$ -module if and only if E_i is a finitely generated (resp., finitely presented) R_i -module for $i = 1, 2$.

Proof of Proposition 2.11. By induction on n , it suffices to prove the assertion for $n = 2$. Since an ideal of $R_1 \amalg R_2$ is of the form $I_1 \amalg I_2$, where I_i is an ideal of R_i for $i = 1, 2$, the conclusion follows easily from Lemma 2.11.

Now, we are able to give a new class of a non-coherent weak coherent rings.

Example 2.13 Let R_1 be a non-coherent weak coherent ring, R_2 be a coherent ring, and $R = R_1 \amalg R_2$. Then:

- 1) R is a weak coherent ring by Theorem 2.10.
- 2) R is not a coherent ring by [5, Theorem 2.4.3]. $\square$

References

- [1] N. Bourbaki, *Commutative algebra, Chapters 1-7*, Springer, Berlin, 1998.
- [2] D. Costa, *Parameterizing families of non-Noetherian rings*, Comm. Algebra 22 (1994), 3997–4011.
- [3] R. Gilmer, *Multiplicative ideal theory*, Pure and Applied Mathematics, No. 12. Marcel Dekker, Inc., New York, 1972.
- [4] S. Glaz, *Finite conductor rings*, Proc. Amer. Math. Soc. 129 (2000), 2833–2843.
- [5] S. Glaz, *Commutative coherent rings*, Lecture Notes in Mathematics, 1371, Springer-Verlag, Berlin, 1989.
- [6] S. Glaz, *Controlling the Zero-Divisors of a Commutative Ring*, Lecture Notes in Pure and Appl. Math., Dekker, 231 (2003), 191–212.
- [7] J.A. Huckaba, *Commutative rings with zero divisors*, Marcel Dekker, New York-Basel, 1988.
- [8] S. Kabbaj, N. Mahdou, *Trivial extensions of local rings and a conjecture of Costa*, Lecture Notes in Pure and Appl. Math., Dekker, 231 (2003), 301–311.
- [9] S. Kabbaj and N. Mahdou, *Trivial extensions defined by coherent-like conditions*, Comm. Algebra 32(10) (2004), 3937–3953.
- [10] N. Mahdou, *On Costa’s conjecture*, Comm. Algebra 29 (2001), 2775–2785.
- [11] N. Mahdou, *On 2-Von Neumann regular rings*, Comm. Algebra 33 (10) (2005) 3489–3496.
- [12] J. J. Rotman, *An Introduction to Homological Algebra*, Academic Press, New York, 1979.